

4. Berkenboom G, Giot C, Unger P, Vachieri JL, Antoine M, LeClerc JL. Plasma endothelin and early coronary endothelial dysfunction in recipients of a cardiac transplant. *Am Heart J*. 1995;129:1178-84.
5. Fedak PW, Rao V, Verma S, Ramzy D, Tumiati L, Miriuka S, et al. Combined endothelial and myocardial protection by endothelin antagonism enhances transplant allograft preservation. *J Thorac Cardiovasc Surg*. 2005;129:407-15.
6. Vassalli G, Gallino A. Endothelial dysfunction and accelerated coronary artery disease in cardiac transplant recipients. Microcirculation Working Group, European Society of Cardiology. *Eur Heart J*. 1997;18:1712-7.
7. Weis M, Wildhirt SM, Schulze C, Rieder G, Wilbert-Lampen U, Wolf WP, et al. Endothelin in coronary endothelial dysfunction early after human heart transplantation. *J Heart Lung Transplant*. 1999;18:1071-9.
8. Dong F, Zhang X, Wold LE, Ren Q, Zhang Z, Ren J. Endothelin-1 enhances oxidative stress, cell proliferation and reduces apoptosis in human umbilical vein endothelial cells: role of ET(B) receptor, NADPH oxidase and caveolin-1. *Br J Pharmacol*. 2005;145:323-33.
9. Wedgwood S, Black SM. Endothelin-1 decreases endothelial NOS expression and activity through ETA receptor-mediated generation of hydrogen peroxide. *Am J Physiol Lung Cell Mol Physiol*. 2005;288:L480-7.
10. Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature*. 1993;362:801-9.
11. Ross R. Atherosclerosis—an inflammatory disease. *N Engl J Med*. 1999;340:115-26.
12. Rifai N, Ridker PM. High-sensitivity C-reactive protein: a novel and promising marker of coronary heart disease. *Clin Chem*. 2001;47:403-11.
13. Ridker PM, Stampfer MJ, Rifai N. Novel risk factors for systemic atherosclerosis: a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *JAMA*. 2001;285:2481-5.
14. Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA*. 2001;286:327-34.
15. Blake GJ, Ridker PM. Inflammatory mechanisms in atherosclerosis: from laboratory evidence to clinical application. *Ital Heart J*. 2001;2:796-800.
16. Rifai N, Buring JE, Lee IM, Manson JE, Ridker PM. Is C-reactive protein specific for vascular disease in women? *Ann Intern Med*. 2002;136:529-33.
17. Albert CM, Ma J, Rifai N, Stampfer MJ, Ridker PM. Prospective study of C-reactive protein, homocysteine, and plasma lipid levels as predictors of sudden cardiac death. *Circulation*. 2002;105:2595-9.
18. Blake GJ, Ridker PM. C-reactive protein and other inflammatory risk markers in acute coronary syndromes. *J Am Coll Cardiol*. 2003;41(4 suppl S):37S-42S.
19. Ridker PM, Morrow DA. C-reactive protein, inflammation, and coronary risk. *Cardiol Clin*. 2003;21:315-25.
20. Libby P, Ridker PM. Inflammation and atherosclerosis: role of C-reactive protein in risk assessment. *Am J Med*. 2004;116(suppl 6A):9S-16S.
21. Verma S, Wang CH, Li SH, Dumont AS, Fedak PW, Badiwala MV, et al. A self-fulfilling prophecy: C-reactive protein attenuates nitric oxide production and inhibits angiogenesis. *Circulation*. 2002;106:913-9.
22. Dlugosz JA, Munk S, Zhou X, Whiteside CI. Endothelin-1-induced mesangial cell contraction involves activation of protein kinase C- $\alpha$ , - $\delta$ , and - $\epsilon$ . *Am J Physiol*. 1998;275(3 Pt 2):F423-32.
23. Park WS, Ko EA, Han J, Kim N, Earm YE. Endothelin-1 acts via protein kinase C to block KATP channels in rabbit coronary and pulmonary arterial smooth muscle cells. *J Cardiovasc Pharmacol*. 2005;45:99-108.
24. Ikonomidis JS, Shirai T, Weisel RD, Derylo B, Rao V, Whiteside CI, et al. Preconditioning cultured human pediatric myocytes requires adenosine and protein kinase C. *Am J Physiol*. 1997;272(3 Pt 2):H1220-30.
25. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976;72:248-54.
26. Matsubara M, Hayashi N, Jing T, Titani K. Regulation of endothelial nitric oxide synthase by protein kinase C. *J Biochem (Tokyo)*. 2003;133:773-81.
27. Partovian C, Zhuang Z, Moodie K, Lin M, Ouchi N, Sessa WC, et al. PKC $\alpha$  activates eNOS and increases arterial blood flow in vivo. *Circ Res*. 2005;97:482-7.
28. Naruse K, Rask-Madsen C, Takahara N, Ha SW, Suzuma K, Way KJ, et al. Activation of vascular protein kinase C- $\beta$  inhibits Akt-dependent endothelial nitric oxide synthase function in obesity-associated insulin resistance. *Diabetes*. 2006;55:691-8.
29. Lee TS, Chao T, Hu KQ, King GL. Endothelin stimulates a sustained 1,2-diacylglycerol increase and protein kinase C activation in bovine aortic smooth muscle cells. *Biochem Biophys Res Commun*. 1989;162:381-6.
30. Lee TS, Hu KQ, Chao T, King GL. Characterization of endothelin receptors and effects of endothelin on diacylglycerol and protein kinase C in retinal capillary pericytes. *Diabetes*. 1989;38:1643-6.
31. McNair LL, Salamanca DA, Khalil RA. Endothelin-1 promotes Ca<sup>2+</sup> antagonist-insensitive coronary smooth muscle contraction via activation of epsilon-protein kinase C. *Hypertension*. 2004;43:897-904.
32. Robin P, Boulven I, Desmyter C, Harbon S, Leiber D. ET-1 stimulates ERK signaling pathway through sequential activation of PKC and Src in rat myometrial cells. *Am J Physiol Cell Physiol*. 2002;283:C251-60.
33. Verma S, Li SH, Badiwala MV, Weisel RD, Fedak PW, Li RK, et al. Endothelin antagonism and interleukin-6 inhibition attenuate the proatherogenic effects of C-reactive protein. *Circulation*. 2002;105:1890-6.
34. Gonon AT, Erbas D, Broijersens A, Valen G, Pernow J. Nitric oxide mediates protective effect of endothelin receptor antagonism during myocardial ischemia and reperfusion. *Am J Physiol Heart Circ Physiol*. 2004;286:H1767-74.
35. Barton M, Haudenschild CC, d'Uscio LV, Shaw S, Munter K, Luscher TF. Endothelin ETA receptor blockade restores NO-mediated endothelial function and inhibits atherosclerosis in apolipoprotein E-deficient mice. *Proc Natl Acad Sci U S A*. 1998;95:14367-72.
36. Li L, Roumeliotis N, Sawamura T, Renier G. C-reactive protein enhances LOX-1 expression in human aortic endothelial cells: relevance of LOX-1 to C-reactive protein-induced endothelial dysfunction. *Circ Res*. 2004;95:877-83.

## Discussion

**Dr Y. Joseph Woo (Philadelphia, Pa).** The authors should be congratulated on a great body of work and, as usual, great graphics from Toronto. Are venous endothelial cells the same as arterial endothelial cells, and are they an adequate surrogate if you are using this as a model for allograft vasculopathy and other sorts of arterial problems?

**Dr Danny Ramzy (Toronto, Canada).** Thank you for the question. We have actually performed experiments looking at that question. Between human saphenous vein endothelial cells, human aortic endothelial cells, and human coronary endothelial cells, once you pass the first passage, and all experiments are performed beyond P1, they all behave the same way as ET-1 and other vasoconstrictors and soluble factors. So in terms of using saphenous vein versus aortic cells, the experiments will yield the same results.

**Dr Mark J. Krasna (Baltimore, Md).** Dr Ramzy, could you tell us a little bit about bosentan? It sounds very exciting. Has it been used clinically in your group in any other arena yet?

**Dr Ramzy.** It has been used clinically. It is approved primarily for primary pulmonary hypertension. It reduces mortality and does reduce pulmonary arterial pressures. It has been used in patients with heart failure, with a few of the studies actually showing harm, likely due to its induction of hypotension. The patient with heart failure died from sudden death, likely from arrhythmia due to

hypotension. So other than in that setting, it has been used safely in patients who can be monitored closely or who have adequate blood pressure. The main problem is the blood pressure.

**Dr Krasna.** Have either of those groups had any relationship between atherosclerosis and bad outcomes and use of bosentan?

**Dr. Ramzy.** It has been looked at mostly in allograft vasculopathy in terms of as a surrogate of atherosclerosis, and it does reduce the burden in terms of amount of disease and the degree of stenosis.

**Dr John G. Byrne (Nashville, Tenn).** What do you think are the principal clinical applications of these findings?

**D. Ramzy.** The clinical applications are severalfold. In terms of atherosclerosis, the reduction of CRP and ET-1 does reduce both the atherosclerotic burden and outcomes such as myocardial infarction. From our results in terms of protein kinase C, we could more specifically target the mechanisms by which CRP results in atherosclerosis and endothelial dysfunction, both in the transplant setting and in atherosclerosis, and especially following transplantation where there is a high degree of vein restenosis, which, if we can modulate basically soluble inflammatory markers like ET-1

and CRP with bosentan or targeted therapy, we could reduce graft restenosis and improve survival and quality of life.

**Dr Byrne.** Are you talking about a vein graft?

**Dr Ramzy.** Vein graft.

**Dr Turki Albacker (Montreal, Canada).** Just for my curiosity, for the clinical application of this study, what do you think about the level of CRP in a patient posttransplant and do you think it is high enough to cause atherosclerosis so you can target it with your therapy? Thanks.

**Dr Ramzy.** There are several studies looking at CRP posttransplant. There are a group of patients who do have elevated levels of CRP, so any level above 3  $\mu\text{g/mL}$  does result in poor outcome. There is actually a recent study that shows any levels above 1.06  $\mu\text{g/mL}$  result in poor outcome. So a slight increase in CRP is able to result in worse outcomes in terms of earlier development of allograft vasculopathy and worse disease, and it has been also shown if you reduce this level, there is a trend toward improvement. Of course, the concentration we used was 25  $\mu\text{g}$ , which is in the intermediate level. We have done studies previous to that. Any level above 3  $\mu\text{g/mL}$  does result in endothelial dysfunction.

### *The Journal of Thoracic and Cardiovascular Surgery* Conflict of Interest Policy

To assure fairness to authors submitting work for consideration in *The Journal of Thoracic and Cardiovascular Surgery*, a mechanism exists for managing conflicts of interest. The editor and each of the section editors complete a "Conflict of Interest" form that identifies any and all relationships with commercial and other academic entities. When the editor has a potential conflict because of a relationship with another entity or author, the editor appoints an alternate editor from among the section editors or editorial board members who assumes the entire responsibility for final decisions on the manuscript in question. The editor does not read the reviews that are submitted nor engage in discussing the manuscript prior to the final decision. When the conflict of interest involves a section editor, a "guest section editor" is appointed who fills the role normally played by the conflicted section editor. All members of the editorial board and reviewers are asked to indicate any conflict of interest when they agree to review a manuscript.